

Commonly Used Post-Op Medications

BLOOD THINNER:

Medication	Dose	Frequency	<u>Comments</u>
ASPIRIN	81 mg	Twice Daily	May cause stomach or nose bleeds

PAIN:

Medication	Dose	Frequency	<u>Comments</u>
ACETAMINOPHEN/CODIENE #3	300/30mg	1 tablet by mouth every 4	• Constipation is a common side effect
(Tylenol #3)		hours AS NEEDED	of this medication
HYDROCODONE (Norco)	5/325 mg	1 tablet by mouth every 4 hours AS NEEDED	 Constipation is a common side effect of this medication This medication contains 325mg acetaminophen per tablet Max dose 4,000 mg/day of acetaminophen total from all sources
TRAMADOL (Ultram)	50 mg (max 400mg/day)	1 tablet by mouth every 8 hours AS NEEDED	 Constipation is a common side effect of this medication May cause confusion, flushing, nausea, & drowsiness

ANTI-INFLAMMATORY:

Medication	<u>Dose</u>	Frequency	<u>Comments</u>
IBUPROFEN	200 mg – 800 mg	3 times daily	DOSAGE MAY VARY
			DO NOT take any other
			NSAIDs
ALEVE	250 mg	1-2 times daily	DO NOT take with any other NSAIDs

NAUSEA:

Medication	Dose	Frequency	<u>Comments</u>
ODANSETRON	4 mg (max dose 16mg/day)	Every 6 hours	May cause headache,
(Zofran)		TAKE ONLY AS NEEDED FOR	constipation, and/or
		NAUSEA	drowsiness
PHENERGAN	25 mg	Every 8 hours	May cause headache,
		TAKE ONLY AS NEEDED FOR	constipation, and/or
		NAUSEA	drowsiness

*Please use over-the-counter stool softeners and/or laxatives to prevent constipation.

* Taking anti-inflammatories along with pain medications will often provide better pain relief.